Concise Review

PROMs Following Root Canal Treatment and Surgical Endodontic Treatment

Jasmine Wong^a, Gary Shun Pan Cheung^{a,b}, Angeline Hui Cheng Lee^a, Colman McGrath^c, Prasanna Neelakantan^{a*}

^a Discipline of Endodontology, Division of Restorative Dental Sciences, Faculty of Dentistry, The University of Hong Kong, Hong Kong SAR, China

^b Department of Dental Surgery, University of Hong Kong – Shenzhen Hospital, Shenzhen, China

^c Dental Public Health, Faculty of Dentistry, The University of Hong Kong, Hong Kong SAR, China

ARTICLE INFO

Article history: Received 24 May 2022 Received in revised form 19 June 2022 Accepted 20 June 2022 Available online xxx

Key words: Minimal important difference Oral health-related quality of life Patient-centered outcome Root canal treatment Surgical endodontics

ABSTRACT

The FDI is currently working on developing a tool to encompass patient-reported outcome measures (PROMs) within the overall assessment of outcomes of endodontic treatment. The outcome of endodontic treatment has traditionally been determined by various clinical and radiographic criteria. However, these parameters do not address the impact of treatment on a patient's oral health-related quality of life (OHRQoL). OHRQoL, a crucial PROM, can be used to understand treatment outcome from a patient-centred perspective, thus improving clinician-patient communication whilst guiding decision-making. This focussed review aims to recount the OHRQoL of patients following nonsurgical root canal treatment and surgical endodontic treatment, with a specific focus on the minimal important difference (MID; the minimum score changes of an outcome instrument for a patient to register a clinically significant change in their OHRQoL and/or oral condition) and the methods used to determine it. The current evidence indicates that the OHRQoL of patients requiring root canal treatment is poorer than those without such need. Accordingly, the literature suggests that OHRQoL improves following nonsurgical or surgical endodontic treatment. However, study methodologies vary widely, and conclusions cannot be drawn with high confidence, nor can MID recommendations be provided. Well-designed clinical studies with baseline measurements and appropriate follow-up time frames are therefore needed. Despite that the literature is rife with outcome studies, research on PROMs is an area that deserves greater attention, particularly in relation to the MID. Determining the MID will facilitate the understanding of changes in outcome scores from the patients' perspective, thus allowing for more informed decision-making in clinical practice.

© 2022 The Authors. Published by Elsevier Inc. on behalf of FDI World Dental Federation. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/)

Introduction

Whilst the goal of root canal treatment is to eliminate infection, relieve pain, restore the health of the periapical tissues, and retain the functionality of the treated tooth,^{1,2} treatment effectiveness and success have traditionally been measured using clinician-reported outcomes that rely on clinical and radiographic criteria.^{1,3} Various criteria for successful root canal treatment have been proposed, most notably the

* Corresponding author. Discipline of Endodontology, Division of Restorative Dental Sciences, Faculty of Dentistry, The University of Hong Kong.

E-mail address: prasanna@hku.hk (P. Neelakantan). https://doi.org/10.1016/j.identj.2022.06.015 "strict"⁴ and "loose"^{5,6} criteria, which are primarily categorised based on complete reversal of the periapical radiolucent area or its arrest. On the other hand, other terms such as favourable, uncertain, and unfavourable¹ as well as healed, healing, nonhealed, and functional⁷ have also been proposed to describe endodontic treatment outcome. Dichotomisation of radiographic appearance as "success" or "failure" to convey prognosis may not be as relevant to patients, as they may have different goals, values, and/or treatment expectations than what the clinician may have in mind.⁸ Evaluating the treatment effectiveness from the patients' perspective, that is, patient-centred outcomes, is of pivotal importance in the context of patient-centred care.⁹ Patient-centred care has been associated with improvements in patient satisfaction

0020-6539/© 2022 The Authors. Published by Elsevier Inc. on behalf of FDI World Dental Federation. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/)

WONG ET AL.

and overall well-being.¹⁰⁻¹² In the context of endodontic therapy, patient-centred care emphasises the elimination of symptoms whilst prioritising functionality.² A recent white paper by the FDI affirms that these treatment philosophies are in line with the concept of "endodontic medicine," which suggests that endodontic diseases should be considered within a greater context, that is, the human body, as they not only affect the health of pulpal and periapical tissues but also impact general health.²

Quality of life (QoL) is one of the key components of patient-centred outcomes that form the basis of the patient -dentist dialogue.¹³ Whilst QoL indicators have been commonly employed throughout health care and general dentistry,¹⁴⁻¹⁶ it has only recently emerged as a topic of interest in endodontics. Current evidence indicates that root canal treatment would positively influence oral health-related QoL (OHRQoL).¹⁷⁻¹⁹ Despite such promising findings, a muchneeded critical appraisal into the potential applications of OHRQoL and different OHRQoL instruments in the field of endodontic research is lacking. Most notably, the minimal important difference (MID) largely remains to be described in detail from the context of root canal treatments.¹⁹ The MID represents the smallest difference in a patient-reported outcome score that is considered clinically significant.²⁰ Thus, to understand whether a change in OHRQoL is meaningful to the patient, determining the MID for the given context is essential. Currently there is a paucity in the evidence pertaining to the MID for OHRQoL measures, demonstrating the need for research in this area.²¹ Therefore, the aim of this review the current evidence to recount the OHRQoL of patients following nonsurgical root canal treatment and surgical endodontics, with a specific focus on MID.

Review

What is OHRQoL, why is it important, and how is it measured?

QoL is "an individual's perception of their position in life in the context of the culture and value systems in which they live and in relation to their goals, expectations, standard and concerns."22 It is a multifaceted construct that incorporates physical, psychological, and social domains. It can be understood in terms of both positive-for example, having the ability to chew and function—and negative—for example, fatigue and pain-dimensions.²² Oral health can have an impact on overall health and QoL by impacting an individual's ability to carry out certain functions, such as chewing, talking, and tasting.²³ Moreover, oral diseases can have psychological and social impacts that can affect an individual's well-being.23,24 Specifically, OHRQoL refers to an individual's self-perceived "comfort when eating, sleeping and engaging in social interaction, their self-esteem and their satisfaction with respect to their oral health."25

Attempts to conceptualise the complex notion of OHRQoL to provide a reference framework for researchers and health care professionals have demonstrated that no one unique dimension can represent OHRQoL (ie, multidimensional) as the different domains work in tandem (ie, integrative).^{14,26,27}

OHRQoL outcome measures are an essential component of patient-centred care since they allow clinicians to holistically evaluate the efficacy of different treatment options in light of the patient's needs and values.¹³ This improves patient-clinician communication and facilitates the treatment decisionmaking process.²⁸ Furthermore, not all patients may have access to the "ideal" care because of social, cultural, and/or economic barriers; hence, patient-centred outcomes can facilitate the setting of individualised treatment goals.²⁸ Where treatments may not be able to eliminate the disease but rather provide palliative/supportive care, improving the QoL may become the primary goal of treatment.²⁹ In regards to public health, patient-centred outcomes may guide the development of health promotion programmes, allocate resources, and evaluate the efficacy of oral health care services.¹³ Furthermore, QoL indicators can be employed in dental research to facilitate evidence-based dentistry, cost-utility analysis, and health service evaluation.³⁰

There are 3 main methods of evaluating OHRQoL: social indicators, global self-ratings, and multiple-item questionnaires.

- a) Social indicators describe community-level social costs of oral disease. Population surveys are carried out to understand the social impact of diminished oral health such as loss of working days, restricted activities, and absence from schools.³¹ However, social indicators provide limited information about the impact of oral health on an individual's OHRQoL.³⁰
- b) Global assessment ratings (global self-ratings or singleitem ratings) involve asking individuals one general question about their oral health status.³² They can be used to determine the responsiveness of an instrument as well as the MID of patient-centred outcomes.³³ This method allows simple and general comparisons; however, it does not adequately reflect the various dimensions of OHRQoL. Therefore, global assessment ratings are often combined with multiple-item questionnaires.³⁴
- c) Multiple-item questionnaires represent the instrument of choice³¹ and can be categorised into generic- and diseaseor condition-specific instruments, such as the Oral Health Impact Profile (OHIP)³⁵ and Geriatric Oral Health Assessment Index,³⁶ respectively.

How has OHRQoL been measured in the endodontic literature?

A literature search was conducted to identify relevant studies using a search strategy that was developed based on previous reviews,¹⁷⁻¹⁹ utilising keywords that related to endodontic treatment and OHRQoL (Table 1). English publications investigating patient-reported outcome measures (PROMs) in relation to endodontic diseases and treatment were identified. After de-duplication and screening of the title and abstract, full texts of the relevant articles were obtained. The references of those articles were then hand searched for any other relevant studies. Hand-searched articles concerning the definition, concepts, and methods of measuring OHRQoL and MID were also included. The first search yielded a total of 449 studies. After removal of

<u>ARTICLE IN PRESS</u>

PROMS IN ENDODONTICS

Table 1 – Search strategy used to identify articles in this narrative review.

Search	Query
#1	"Root canal treatment" OR "root canal therapy" OR "endodontic treatment" OR "endodontics" OR "root canal retreatment" OR
#2	"Endodontic retreatment" "Apicoectomy" OR "apicectomy" OR "periradicular sur- gery" OR "Endodontic surgery" OR "apical surgery" OR "periapical surgery" OR "root-end surgery" OR "root-end vocation"
#3	"Patient-reported outcome measures" OR "health- related quality of life" OR "oral health-related quality of life" OR "Quality of life" OR "quality of life index" OR "patient satisfaction" OR "general quality of life" OR "WHOQoL" OR "QoL" OR "health utility index" OR "SF-36" OR "SF-12" OR "SF-9" OR "SF-6" OR "EUROQoL" OR "EQ-5D"
#4 #5	#1 OR #2 #3 AND #4

duplicates and hand searching of references, 32 clinical studies and 3 systematic reviews were identified. Tables 2 through 4 summarise the methodologic characteristics and key findings of the OHRQoL studies identified in the literature related to nonsurgical root canal treatment and surgical endodontics.

Dugas et al³⁷ conducted the first study investigating the OHRQoL of endodontic patients and, since then, the importance of PROMs has been thrust into the limelight, prompting a steady growth of OHRQoL studies in relation to endodontic disease and treatment. The ideal instrument for PROMs should be appropriate, reliable, valid, responsive, and interpretable.^{9,31} However, as a "goldstandard" instrument for endodontic patients remains to be established, a myriad of measures has been employed to characterise the impact of root canal treatment on OHRQoL.¹⁹ Currently, the most frequently utilised instrument in the endodontic literature is the OHIP, specifically the OHIP-14.

The OHIP-14, which is a shortened version of the original OHIP-49, was developed based on Locker's conceptual model of oral health.²⁶ The questionnaire is subdivided into 7 domains: functional limitation, physical pain, psychological discomfort, physical disability, social disability, and handicap. The patient answers based on how often they have encountered each scenario within a specific time frame, usually 12 months, using a 5-point Likert scale. The scores are summated, with a higher total score indicating poorer levels of OHRQoL.³⁸ Other variations of the OHIP have been found in the endodontic literature as well, such as the OHIP-17³⁷ and the OHIP-14_sev.³⁹

A key benefit of using the OHIP-14 in the context of endodontics is that it has been confirmed to be sensitive enough to detect changes in patients' OHRQoL following endodontic treatment.⁴⁰ It also has been translated and validated in multiple languages,⁴¹⁻⁴³ allowing adaptability for different cultural contexts. However, there exists much variation on how researchers interpreted the outcomes from OHIP-14. Some have dichotomised the results into "no impact" and "impact,"⁴⁴ whilst others defined poor OHRQoL as scores that were amongst the upper quartile of the study group.⁴⁰ Furthermore, how studies deduced improvement in OHQRoL was not standardised in the endodontic literature, with some inferring it from changes in the total score whilst others based it on changes to the individual domains or even the individual item level.¹⁹ It has been suggested that summed scores and domain-level analysis are favoured over itemlevel analysis.⁴⁵

Other OHRQoL instruments have made also appeared in the endodontic literature, such as the Patient Perception Questionnaire,⁴⁶⁻⁵³ Health-related QoL Index,⁵⁴ Oral Impact on Daily Performance (OIDP),55 General Oral Health Assessment Index,⁵⁶ OHRQoL instrument,⁵⁷ Post-operative QoL questionnaire, 58,59 OHRQoL research tool,60 and the QoL Scale.⁶¹ Although these instruments may provide an alternative means to measure PROMs, there are several factors that may hinder their widespread use in endodontic research. First, the responsiveness of most of these instruments have not been thoroughly investigated regarding OHRQoL changes associated with endodontic disease and treatment. Second, given their limited use throughout the endodontic literature, comparisons between studies may be challenging, which could prevent an accurate quantitative synthesis (ie, metaanalyses).

Health-related QoL measures, such as the General Health Questionnaire⁶² and the EuroQoL-5D-5L instrument,⁶³ are sometimes utilised to provide an additional assessment of the patients' general QoL. These instruments may allow researchers to evaluate how endodontic treatment–related factors can affect a patients' self-perceived general health and overall well-being. However, the sensitivity of generic questionnaires is known to be inferior to disease-specific questionnaires.⁶⁴

Apart from the choice of instrument, a crucial element for consideration is the time period of assessment. Ideally, a baseline measurement of the patients' OHRQoL must be provided. Cross-sectional studies only capture the OHRQoL at a single time point, generally months to years posttreatment, which may result in susceptibility to recall bias. Prospective studies and randomised clinical trials thus possess a clear advantage. However, significant variation exists in regard to the evaluation periods. Whilst several studies have applied extended evaluation periods, for example, 1 year^{54,65} to 2 years,^{66,67} others reported postoperative assessments of only up to 7 days, sometimes without any preoperative baseline measurement.46-52 It has been suggested that limited time frames, for example, 6 months or less, are insufficient to evaluate changes in OHRQoL as they are limited to describing the initial posttreatment recovery. Hence, follow-up periods of approximately 1 year may be more suitable.^{18,45} On the other hand, further lengthening the period of evaluation may result OHRQoL fluctuations due to other oral diseases having emerged.⁵⁴

Study	Type of study; country	Sample	Intervention	Time frame	OHRQoL measure	MID determined	Key findings
Dugas et al 2002 ³⁷	Cross-sectional study; Canada	119 patients	Root canal treatment	Treatment within 2 years from the begin- ning of the study	OHIP-17 on the impact of dis- ease with cor- responding questions on the impact of treatment	No	OHRQoL (OHIP-17 score) was most impacted by the disease in the domains of "physical pain" and "psychological disability." Improvement after root canal treatment was experienced in all aspects of OHRQoL (OHIP-17 items). Patients with "painful aching" preoperatively had the largest rate of improvement. There were significant differences in the improvement of various aspects of OHRQoL associated with different factors (operator experi- ence, PAI score, education, missing teeth).
Jordan et al 2009 ⁵⁴	Prospective study; Republic of Gambia	15 patients	Root canal treat- ment: basic treatment protocol	Before treatment and 1 day, 5 days, 6 months, 12 months after treatment	HRQoL index	No	HRQoL improved (HRQoL index score decreased) with time after treatment, particularly in relation to pain, chewing ability and ability to work. The largest improvement was seen immediately after treatment (ie, 1 day). HRQoL index score fluctuated at 6 and 12 months.
Wright et al 2009 ⁵⁷	Prospective study; USA	63 patients (15 endodontic, 16 denture, 32 recall) at base- line; 44 patients at fol- low-up	Root canal treat- ment, denture replacement, or recall with no apparent disease	Before treatment and 3 months after treatment	6- and 12- item OQOL instru- ment, global self-report of oral health	No	No significant difference of OHRQoL (OQOL scores) between groups at both time points. OHRQoL improved (OQOL instrument scores decreased) after treatment, with small effect sizes in the end- odontic and recall group and moderate effect sizes in the denture group.
Gatten et al 2011 ⁶⁹	Cross-sectional study; USA	37 patients (17 endodontic, 20 implant)	Root canal treat- ment vs implant treatment	After treatment, no specific time frame stated	OHIP-14, focus group discussions	No	The majority of patients did not experience any impact on OHRQoL (OHIP-14 score) after treat- ment. There was no significant difference of OHR- QoL between groups. OHRQoL was most impacted in the domains of "physical pain" and "psychological disability" in both groups. The end- odontic group experienced significantly higher scores in the domains "psychological discomfort" and "psychological disability" compared to the implant group. Most participants expressed a desire to retain their natural dentition when possible.
Yu et al 2012 ⁵⁵	Cross-sectional study; Singapore	127 patients with 185 persistent lesions	Root canal treat- ment with per- sistent end- odontic lesion and painful exacerbations	After treatment, no specific time frame stated	Modified OIDP	No	Out of the patients who experienced painful epi- sodes, a large proportion reported no to very minor effect on their daily living.

Table 2 – Summary of study characteristics involving nonsurgical root canal treatment and retreatment.

WONG ET AL.

4

(continued on next page)

Table 2 (Continued)

Study	Type of study; country	Sample	Intervention	Time frame	OHRQoL measure	MID determined	Key findings
Liu et al 2012 ⁶²	Case-control study; Hong Kong	200 patients (100 endodontic patients, 100 control patients, ie, periodontal maintenance)	Indicated for root canal treat- ment vs peri- odontal maintenance	Before treatment or scheduled for periodontal maintenance	OHIP-14 Chinese version, GHQ- 12 Chinese version	No	OHRQoL (OHIP-14 score) and psychological well- being (GHQ-12 score) were poorer in the endodon- tic patient group compared to the periodontal maintenance group.
Liu et al 2014 ⁷²	Cross-sectional study; Hong Kong	412 patients	Indicated for root canal treatment	Before treatment	OHIP-14 Chinese version	No	OHRQoL (OHIP-14 scores) was poorer in endodontic patients and was associated with multiple teeth needing treatment, older age, and increased pain.
Liu et al 2014 ⁴⁰	Prospective study; Hong Kong	253 patients at 1- month recall; 213 patients at 6-month recall	Root canal treatment	Before treatment and 1 month and 6 months after treatment	OHIP-14 Chinese version, global item rating of oral health improvement	No	OHRQoL significantly improved after treatment (OHIP-14 scores decreased) at both 1-month and 6- month recalls, with moderate and large effect sizes respectively. Self-ratings of improvement in oral health were significantly associated with changes in OHRQoL (OHIP-14 scores) and PAI scores.
Vena et al 2014 ⁷¹	Cross-sectional study; USA	1257 patients	Root canal treatment	Treatment within the last 3-5 years	OHIP-14	No	"Pain upon percussion" and "periapical pathosis" were associated with a negative impact on OHR- QoL (OHIP-14 scores).
Montero et al 2015 ³⁹	Prospective study (cross-sectional OHRQoL com- ponent); Spain	250 patients	Root canal treatment	Before treatment	OHIP-14_sev Spanish version	No	OHRQoL (OHIP-14_sev score) was most impacted at baseline in the domains of "physical pain" and "psychological discomfort." There were significant differences in OHRQoL domains associated with various factors (tooth type, socioeconomic status, age, gender).
He et al 2017 ⁶⁷	Prospective study; USA	52 patients	Root canal retreatment	At entry (before treatment) and 1 week, 1 month, 6 months, 12 months, and 24 months after treatment	OHIP-17	No	OHRQoL improved (OHIP-17 scores decreased) signif- icantly after root canal retreatment. The largest improvement occurred within the first week, after which the improvement rate slowed.
Hamasha & Hatiwsh 2017 ⁶⁸	Prospective study; Jordan	302 patients (101 were treated by 22 undergradu- ate students, 100 were treated by four graduate stu- dents and 101 participants were treated by three endodon- tic specialists)	Root canal treatment	Before treatment and 2 weeks after treatment	OHIP-17 in Arabic version	No	The median impact of pulpal disease on OHRQoL (OHIP-17 score) was low overall. The highest impact was observed in the domains "physical pain" and "psychological disability." OHRQoL improved (OHIP-17 scores decreased) after treat- ment. No significant difference in improvement associated with operator level. There were signifi- cant differences in OHRQoL associated with vari- ous factors (presence of gingival inflammation, history of missing teeth, pulp status).

Table 2 (Continued)

Study	Type of study; country	Sample	Intervention	Time frame	OHRQoL measure	MID determined	Key findings
Chew et al 2019 ⁶⁶	Prospective study; Australia	1096 patients at baseline, 438 at 2-year recall	Root canal treat- ment vs other dental services (extraction, restorations, prosthodontics, periodontics, preventative treatment, and scale and clean)	Baseline and 2 years	OHIP-14, global transition statement of change	No	Root canal treatment group had significantly lower odds for good/improved OHRQoL outcomes (lower OHIP-14 and GTSC scores) at the 2-year review compared to all dental services, but not individual treatment groups. The "preventative" and "scale and clean" groups had significantly higher odds for improved health.
Iqbal et al 2020 ⁶⁰	Cross-sectional study; Pakistan	57 patients	Root canal treatment	After treatment, no specific time frame stated	OHRQoL research tool	No	A majority of patients expressed no impact on their OHRQoL after treatment in all 4 domains (physical function, psychological, social, and pain). A mod- erately good level of OHRQoL was observed amongst patients receiving root canal treatment. There were significant differences in the improve- ment of OHRQoL in the domains of physical func- tion, psychological and pain associated with marital status, smoking status, and gender, respectivelv.
Wigsten et al 2020 ⁶³	Prospective study: Sweden	85 patients (48 extraction, 37 endodontic)	Root canal treat- ment vs extraction	Baseline (at the initiation of treatment) and 1-month follow up	OHIP-14 and EQ- 5D-5L Swedish versions	No	No significant difference of OHRQoL (OHIP-14 scores) between time points for both groups. The extrac- tion group registered greater "embarrassment" compared to the endodontic group. HRQoL (EQ-5D- 5L score) was significantly improved in the end- odontic group only.

EQ-5D-5L, EuroQoL-5D-5L; GHQ-12, general health questionnaire-12; GTSC, global transition statement of change; HRQoL, health-related quality of life; MID, minimal important difference; OHIP-14, Oral Health Impact Profile-14; OHIP-17, Oral Health Impact Profile-17; OHIP-14_sev, Oral Health Impact Profile-14 severity; OIDP, Oral Impact on Daily Performance; OHRQoL, oral health-related quality of life; OQOL, Oral Health-Related Quality of Life instrument; PAI, periapical index.

WONG ET AL.

Study	Type of study; country	Sample size	Intervention	Time frame	OHRQoL measure	MID determined	Key findings
Pasqualini et al 2016 ⁵⁸	Randomised clinical trial; Italy	47 patients (23 rotary, 24 reciprocating)	Reciprocating instru- mentation vs rotary instrumentation	Every day from the day of surgery to 7 days postoperatively	POQoL questionnaire	No	OHRQoL improved with time (POQoL score decreased). Per- ceived OHRQoL was significantly better for the rotary group.
Bartols et al 2016 ⁷³	Prospective study; Germany	137 patients (71 reciprocating group, 66 hand file group)	Reciprocating instru- mentation vs hand instrumentation	In the week before treatment and in the week before completion of treatment (ie, 14 days after initial treatment)	OHIP-14 German version	No	OHRQoL improved (OHIP-14 scores decreased) significantly after root canal treatment. No significant difference of OHRQoL (OHIP-14 scores) between groups.
Yaylali et al 2017 ⁶¹	Randomised con- trolled trial; Turkey	70 (35 with foramen enlargement, 35 without)	Foraminal enlarge- ment vs no forami- nal enlargement	Every day from the day of surgery to 7 days postoperatively	QoLS	No	No significant difference of OHR- QoL (QoLS scores) between groups.
Oliveira et al 2018 ⁷⁴	Randomised clinical trial; Brazil	58 patients (29 recip- rocating, 29 rotary)	Reciprocating instru- mentation vs rotary instrumentation	24 hours after treatment	OHIP-14 Brazilian version	No	No significant difference of OHR- QoL (OHIP-14 scores) between groups. Higher postoperative pain (VAS score) was associated with poorer OHRQoL (higher OHIP-14 score).
Yavari et al 2019 ⁵⁹	Randomised clinical trial; Iran	196 patients (64 dexamethasone, 66 betamethasone, 64 saline)	Local infiltration of betamethasone vs dexamethasone vs saline after 1-visit root canal treatment	Before treatment; 6, 12, 24, 48, and 72 hours after treatment; and 7 days after treatment	POQoL questionnaire	No	Both the corticosteroid groups had significantly higher OHRQoL (POQoL scores) than the placebo group. A decrease in pain was associated with an increased in OHRQoL (POQoL scores).
Diniz-de-Fig- ueiredo et al 2020 ⁶⁵	Randomised con- trolled trial; Brazil	88 patients at 6 months (46 man- ual group, 42 recip- rocating group); 87 patients at 12 months (42 man- ual group, 45 recip- rocating group)	Reciprocating instru- mentation and sin- gle cone obturation vs hand file instrumenta- tion and lateral compaction obturation	Prior to treatment and 6 and 12 months after treatment	OHIP-14 Brazilian version	Yes	OHRQoL improved (OHIP-14 scores decreased) significantly after root canal treatment with moderate to large effect sizes. At 6 months, the manual protocol was associ- ated with poorer OHIP-14 scores. At 12 months, there was no sig- nificant difference between groups. There was no significant difference between 6-month and 12-month OHIP-14 scores within groups.

Table 3 - Summary of study characteristics involving procedural aspects of the nonsurgical root canal treatment protocol.

MID, minimal important difference; OHIP-14, Oral Health Impact Profile-14; OHRQoL, oral health-related quality of life; POQoL, Postoperative Quality of Life; QoLS, Quality of Life Scale; VAS, visual analogue scale.

Study	Type of study; country	Sample size	Intervention	Time	OHRQoL measure	MID determined	Key findings
Tsesis et al 2005 ⁵³	Prospective study; Israel	63 patients (31 tradi- tional surgery, 32 microsurgery)	Surgical endodontic treatment with traditional vs microsurgical techniques	Every day from the day of surgery to 7 days postoperatively	PPQ	No	The traditional surgery group has significantly more pain than the microsurgery group. The microsurgery group had significantly more diffi- culty in mouth opening, speaking, and mastication than the traditional sur- gery group.
Del Fabbro et al 2009 ⁴⁸	Randomised clin- ical trial; Italy	38 patients (19 PBI, 19 SI)	Surgical endodontic treatment with PBI vs SI flap designs	Every day from the day of surgery to 7 days postoperatively	PPQ	No	The PBI group experienced significantly faster reduc- tion in pain compared, less swelling, and less chewing impairment than the SI group.
Del Fabbro et al 2012 ⁴⁷	Randomised clin- ical trial; Italy	36 patients (18 con- trol, 18 PRGF)	Surgical endodontic treatment with PRGF vs none	Every day from the day of surgery to 7 days postoperatively	PPQ	No	The PRGF group reported significantly less pain and swelling, less consump- tion of analgesics, and improved functional activities compared to the control group.
Taschieri et al 2014 ⁵¹	Retrospective study; Italy	20 patients (12 con- trol, 8 PRGF)	Sinus perforation management with PRGF vs none dur- ing surgical end- odontic treatment	Every day from the day of surgery to 7 days postoperatively	PPQ	No	The PRGF group reported significantly better OHR- QoL in multiple domains (eg, swelling, bad breath/ taste, pain, various func- tional activities) com- pared to the test group.
Meschi et al 2018 ⁴⁹	Randomised con- trolled trial; Belgium	50 patients (25 LPRF, 25 control)	Surgical endodontic treatment with LPRF vs none	Every day from the day of surgery to 7 days postoperatively	PPQ	No	No significant difference of OHRQoL (patient per- ceived postoperative symptoms) between groups.
Metin et al 2018 ⁵⁶	Prospective study; Turkey	71 (34 LLLT, 37 control)	Surgical endodontic treatment with LLLT vs none	1, 3, and 7 days postoperatively	GOHAI and OHIP- 14 Turkish versions	No	The LLLT group reported significantly better OHR- QoL (OHIP-14 and GOHAI scores) compared to the

Table 4 – Summary of study characteristics involving surgical endodontic treatment.

(continued on next page)

control on day 1 and 3 postoperatively.

WONG ET AL.

00

Study	Type of study; country	Sample size	Intervention	Time	OHRQoL measure	MID determined	Key findings
Soto-Peñaloza et al 2020 ⁵⁰	Randomised clin- ical trial; Spain	50 patients (25 A-PRF +, 25 control)	Surgical endodontic treatment with A- PRF+ vs none	Every day from the day of surgery to 7 days postoperatively	PPQ	No	The A-PRF+ group reported significantly better speech and sleep functions com- pared to the test group.
Khoo et al 2020 ⁴⁴	Cross-sectional study; Singapore	150 patients (75 retreatment, 75 apical surgery)	Root canal retreat- ment vs surgical endodontic treatment	6 to 24 months after treatment	OHIP-14 Chinese and Malay ver- sions	No	Impact on OHRQoL (OHIP- 14 scores) was low, with no significant difference between groups. Impact most commonly experi- enced in the domains of "physical pain" and "psychological dis- comfort." Poorer OHRQoL (higher OHIP-14 scores) associated with women and presence of preopera- tive pain. There was no correlation between OHIP-14 sores and healing outcome.
Tuk et al 2021 ⁵²	Prospective study; Amsterdam	133 patients	Surgical endodontic treatment	Every day from the day of surgery to 7 days postoperatively	OHIP-14 Dutch version supple- mented with questions on postoperative symptoms	No	OHRQoL generally improved throughout the week (OHIP-14 score decreased). There were significant differences in OHRQoL associated with various factors (age, post- operative infection, smoker).
Bharathi et al 2021 ⁴⁶	Randomised clin- ical trial; India	40 patients (20 piezo- surgery, 20 control)	Surgical endodontic treatment with piezosurgery pro- tocol vs conventional	Every day from the day of surgery to 7 days postoperatively	PPQ	No	The piezosurgery group experienced significantly less swelling and less pain compared to the control group.

A-PRF+, advanced platelet-rich fibrin; GOHAI, General Oral Health Assessment Index; LLLT, low-level laser therapy; LPRF, leukocyte- and platelet-rich fibrin; MID, minimal important difference; OHIP-14, Oral Health Impact Profile-14; OHRQoL, oral health–related quality of life; PBI, papilla-based incision; PPQ, Patient Perception Questionnaire; PRGF, plasma rich in growth factors; SI, sulcular incision.

OHRQoL associated with endodontic disease and treatment

Endodontic diseases have been found to negatively impact OHRQoL^{62,68} particularly in the domains of physical pain, psychological disability, and psychological discomfort. 37,39,44,68,69 Studies have reported OHRQoL improvement after primary^{37,40,57,66} and secondary⁶⁷ nonsurgical root canal treatment as well as surgical endodontic treatment.52 Conversely, studies have also reported no significant difference in OHRQoL after nonsurgical treatment.60,63 These contrasting findings may be explained by the heterogeneity of endodontic patients in the disease- (ie, preoperative symptoms), treatment- (ie, complications), and patient-related factors (ie, experience of the treatment, psychosocial factors, and patient values). Furthermore, some endodontic diseases may manifest as "painless" ailments⁷⁰ resulting in minimal perceived impact on OHRQoL.44,68 Therefore, it is likely that the extent of impact also depends on the severity of the symptoms, functional limitation, and psychosocial impairment. Nevertheless, based on the available literature, it may be considered that endodontic treatment generally improves the OHRQoL.17-19

When root canal treatment was compared with other dental services such as extraction, restoration, prosthodontic, periodontal, and preventative treatment, there were no differences when compared to individual treatment groups.⁶⁶ Similarly, no significant difference was found in the OHRQoL between patients who had received root canal treatment vs extraction, although those in the extraction group expressed higher levels of embarrassment.⁵⁷ It has also been reported that a consistent theme with most patients was the desire to keep their natural dentition.⁶⁹

Factors that may influence the OHRQoL of endodontic patients

A large cross-sectional study identified 3 key factors that were associated with poorer OHRQoL: multiple teeth needing treatment, retreatment, and pain.⁶² Both preoperative pain^{44,62} and persistent pain following treatment^{55,71} were found to negatively impact PROMs. The association between OHRQoL and different sociodemographic factors such as gender, age, socioeconomic status, and marital status and has been demonstrated in some studies^{39,44,52} and refuted in others.^{68,72} Similarly, studies on the impact of operator experience have reported conflicting findings, although patient satisfaction was consistently higher when treated by specialists.^{37,68}

Multiple clinical studies have investigated how various procedural aspects of root canal treatment may impact OHR-QoL. This includes local infiltration of corticosteroids,⁵⁹ different instrumentation protocols,^{58,65,73,74} obturation techniques,⁶⁵ and extent of foraminal enlargement.⁶¹ In terms of surgical endodontics, the use of microsurgical protocols,⁵³ peizosurgery instruments,⁴⁶ papilla-based flap designs,⁴⁸ low-level laser therapy,⁵⁶ and autologous platelet concentrates have also been evaluated in the context of PROMs. Except for the study by Diniz-de-Figueiredo et al,⁶⁵ the period of evaluation was relatively short, spanning 2 weeks at most. Again, short evaluation times may be insufficient to thoroughly assess OHRQoL beyond patients' initial recovery.⁴⁵ Although it is entirely conceivable that different

procedural aspects can impact the immediate postoperative experience of the patient, the influence of these factors on the long-term transformation of OHRQoL remains questionable.

A potential relationship between OHRQoL and clinical outcome has been implicated but not well substantiated in several studies.^{37,40} One study found an association between poorer OHRQoL and patients who had an endodontically treated tooth with persistent disease.³⁷ The authors, however, encouraged caution in the interpretation of these results as radiographic outcomes given that this was a cross-sectional survey.³⁷ Another study reported that all domains of OHIP-14 were significantly associated with self-perceived improvement in oral health, whilst some domains changed with respect to improvement in radiographic outcome.⁴⁰ In general, clinical and radiographic parameters of success do not always reflect the changes in OHRQoL, whilst subjective measures such as self-perceived oral health appear to show a stronger correlation/association.^{40,75,76}

MID: a critical element for future research

The extent of benefit gained from any treatment is important for all stakeholders (eg, clinicians, patients, policymakers) to make changes in treatment philosophies. From the context of PROMs, the magnitude of change is a crucial element that represents the benefit gained from treatment. Statistical methods such as calculating the effect size and half of the standard deviation have also been utilised to indicate the magnitude of change.^{40,57,65} Global statements of change are widely used to assess the patients' self-perceived change in oral health status.^{40,57,66} These methods can also be used to infer the responsiveness of the OHRQoL instrument.

The concept of responsiveness was first introduced by Guyatt et al⁷⁷ and was used to describe the ability of an instrument measuring patient-centred outcomes to detect a clinically important change. Subsequently, Jaeschke et al⁷⁸ suggested the term minimal clinical important difference to denote the smallest difference in score which patients perceive as being beneficial. Since then, a myriad of terms have been introduced to represent similar concepts, for example, MID,²⁰ minimally important change,⁷⁹ subjectively significant difference,⁸⁰ and clinical important difference.⁸¹ Despite the many variations in terminology, it has been suggested that MID is the term that is generally used in the literature.⁸²

Ascertaining the MID of PROMs provides multiple benefits.¹⁹ Interpreting the changes in the outcome scores remains unintuitive to both the clinicians and the researchers because statistical significant differences do not reflect the inherent value of the change in score to the patient. Thus, determining the MID allows health care professionals and researchers to interpret the significance of the changes in outcome score.²¹ Furthermore, improvement or deterioration in clinical measurements does not always align or adequately represent the changes from the patient's perspective. Therefore, the MID facilitates better understanding of a patient's self-perceived changes in oral health status and OHRQoL.³³

There are 2 main methods used to determine the MID: anchor-based methods and distribution-based methods. Anchor-based methods use an external marker of change,

<u>ARTICLE IN PRESS</u>

PROMS IN ENDODONTICS

that is, the anchor, to identify whether the difference in outcome score is of clinical significance.^{82,83} The anchor can be objective or subjective; however, the latter is more widely used and is often operationalised in the form of a global statement of change.^{82,84} Distribution-based methods make inferences from the data collected from the patient-reported outcome instrument whilst using the distribution of the scores to calculate the MID value.83 These statistical approaches most commonly include the calculation of effect size, standard error of measurement, and ratios of standard deviation.⁸⁵ The major benefit of using distribution-based methods is that no additional data are required.⁸² However, many argue that the MID of PROMs can only truly be assessed through an understanding of the patient's subjective experience. Hence, it has been suggested that different approaches should be combined to determine MID values, with distribution-based methods providing a supporting role whilst anchor-based methods provide primary evidence.^{84,86,87}

MID has been thoroughly investigated in regards to various medical conditions and treatments.77,88-92 On the other hand, its appearance in OHRQoL research has been lacking.²¹ The majority of studies have solely used distributional methods.93-95 One of the first studies to use an anchor-based approach reported that the MID for OHIP-14 was 5 scale points for an elderly dental population.³² When applied to a group of periodontal patients, the MID was around 5 scale points for the OIDP index.⁹⁶ A recent study investigating OHR-QoL after oral rehabilitative treatment reported a range of values for variants of the OHIP, including 14 scale points for the OHIP-49 and 3 scale points for the OHIP-14.97 Only one study has investigated the MID of OHRQoL for endodontic patients; however, only distribution-based approaches were utilised.65 To the best of our knowledge, there are no studies evaluating MID of OHRQoL for endodontic patients using anchor-based methods. It has been emphasised that specific MID values should be interpreted within the context of a given application, with special attention paid to the OHRQoL instrument used and the characteristics of the study group.⁸⁶

Given that the FDI is currently working on the development of an oral health measurement tool, which incoprorates patient-centred outcomes as a measure in the assessment of oral health outcomes,² research on MID may significantly improve the clinical usefulness of such tools.

Conclusions

A combination of PROMs with clinical and radiographic outcome measures can result in a more comprehensive understanding of the impact of endodontic treatment and the value of different treatment modalities. The evidence supports that endodontic diseases can have a negative impact on OHRQoL, and whilst endodontic treatment has been shown to enhance patients' OHRQoL, the extent of improvements vary. To strengthen the current evidence, well-designed large-scale clinical studies are needed to determine the effect of root canal treatment on OHRQoL in comparison with alternative modes of treatment, such as extraction and/or implants. These studies should include a baseline measurement of OHRQoL, a suitable time period of assessment, and an appropriate choice of instrument. In addition, there is a need to develop endodontic-specific OHRQoL instruments to be used in tandem with generic OHRQoL instruments in future research. Last, investigating the MID is elemental for a thorough interpretation of OHRQoL measures. Understanding the MID gives insight into both the magnitude and value of change after an intervention from the patient's perspective and hence should be a prime focus of future studies.

Author contributions

Conceptualisation: Jasmine Wong, Gary Shun Pan Cheung, Prasanna Neelakantan

Literature search and initial review: Jasmine Wong, Prasanna Neelakantan

Writing–first draft: Jasmine Wong, Gary Shun Pan Cheung, Prasanna Neelakantan, Angeline Lee, Colman McGrath

Writing—review and final version: Jasmine Wong, Gary Shun Pan Cheung, Prasanna Neelakantan, Angeline Lee, Colman McGrath

Funding acquisition: Gary Shun Pan Cheung, Prasanna Neelakantan

Conflict of interest

The authors declare that they have no conflicts of interest.

REFERENCES

- European Society of Endodontology. Quality guidelines for endodontic treatment: consensus report of the European Society of Endodontology. Int Endod J 2006;39(12):921–30.
- Peters OA, Seeberger GK. White paper on endodontic care 2019. FDI World Dental Federation; 2019. Available from: https://www.fdiworlddental.org/resource/white-paper-endodontic-care Accessed 23 May 2022.
- Orstavik D, Kerekes K, Eriksen HM. The periapical index: a scoring system for radiographic assessment of apical periodontitis. Endod Dent Traumatol 1986;2(1):20–34.
- Strindberg LZ. The dependence of the results of pulp therapy on certain factors. An analytical study based on radiographic and clinical follow-up examination. Acta Odontol Scand 1956;14:1–174.
- Bender IB, Seltzer S, Soltanoff W. Endodontic success a reappraisal of criteria. II. Oral Surg Oral Med Oral Pathol 1966;22 (6):790–802.
- Bender IB, Seltzer S, Soltanoff W. Endodontic success a reappraisal of criteria. I. Oral Surg Oral Med Oral Pathol 1966;22 (6):780–9.
- American Association of Endodontists. Glossary of endodontic terms. Available from: https://www.aae.org/specialty/ clinical-resources/glossary-endodontic-terms/ Accessed 20 January 2022.
- Friedman S, Mor C. The success of endodontic therapy-healing and functionality. J Calif Dent Assoc 2004;32(6):493–503.
- 9. Fitzpatrick R, Davey C, Buxton MJ, Jones DR. Evaluating patient-based outcome measures for use in clinical trials. Health Technol Assess 1998;2(14):i-iv 1–74.
- 10. Newsome PR, McGrath C. Patient-centred measures in dental practice: 1. An overview. Dent Update 2006;33(10):596–8 600.

- Scambler S, Asimakopoulou K. A model of patient-centred care - turning good care into patient-centred care. Br Dent J 2014;217(5):225–8.
- Rathert C, Wyrwich MD, Boren SA. Patient-centered care and outcomes: a systematic review of the literature. Med Care Res Rev 2013;70(4):351–79.
- McGrath C, Bedi R. The value and use of 'quality of life' measures in the primary dental care setting. Prim Dent Care 1999;6 (2):53–7.
- Chen MS, Hunter P. Oral health and quality of life in New Zealand: a social perspective. Soc Sci Med 1996;43(8):1213–22.
- Feu D, de Oliveira BH, de Oliveira Almeida MA, Kiyak HA, Miguel JA. Oral health-related quality of life and orthodontic treatment seeking. Am J Orthod Dentofacial Orthop 2010;138 (2):152–9.
- **16.** Malele-Kolisa Y, Yengopal V, Igumbor J, Nqcobo CB, Ralephenya TRD. Systematic review of factors influencing oral health-related quality of life in children in Africa. Afr J Prim Health Care Fam Med 2019;11(1):e1–e12.
- Antunes LS, Souza CR, Salles AG, Gomes CC, Antunes LA. Does conventional endodontic treatment impact oral healthrelated quality of life? A systematic review. Eur Endod J 2018;3 (1):2–8.
- Leong DJX, Yap AU. Quality of life of patients with endodontically treated teeth: a systematic review. Aust Endod J 2020;46 (1):130–9.
- Neelakantan P, Liu P, Dummer PMH, McGrath C. Oral healthrelated quality of life (OHRQoL) before and after endodontic treatment: a systematic review. Clin Oral Investig 2020;24 (1):25–36.
- Guyatt GH, Osoba D, Wu AW, Wyrwich KW, Norman GR. Methods for explaining the clinical significance of health status measures. Mayo Clin Proc 2002;77(4):371–83.
- Masood M, Masood Y, Saub R, Newton JT. Need of minimal important difference for oral health-related quality of life measures. J Public Health Dent 2014;74(1):13–20.
- WHOQOL. The World Health Organization quality of life assessment (WHOQOL): position paper from the World Health Organization. Soc Sci Med 1995;41(10):1403–9.
- 23. Petersen PE. The World Oral Health Report 2003: continuous improvement of oral health in the 21st century—the approach of the WHO Global Oral Health Programme. Community Dent Oral Epidemiol 2003;31(Suppl 1):3–23.
- 24. Reisine ST. Dental health and public policy: the social impact of dental disease. Am J Public Health 1985;75(1):27–30.
- 25. US Department of Health and Human Services. Oral health in America: a report of the surgeon general. Available from: https://www.nidcr.nih.gov/sites/default/files/2017-10/ hck1ocv.%40www.surgeon.fullrpt.pdf Accessed 20 February 2022.
- **26.** Locker D. Measuring oral health: a conceptual framework. Community Dent Health 1988;5(1):3–18.
- Wilson IB, Cleary PD. Linking clinical variables with healthrelated quality of life. A conceptual model of patient outcomes. JAMA 1995;273(1):59–65.
- Sischo L, Broder HL. Oral health-related quality of life: what, why, how, and future implications. J Dent Res 2011;90 (11):1264–70.
- Najman JM, Levine S. Evaluating the impact of medical care and technologies on the quality of life: a review and a critique. Soc Sci Med 1981;15F(2–3):107–15.
- **30.** Allen PF. Assessment of oral health related quality of life. Health Qual Life Outcomes 2003;1:40.
- Bennadi D, Reddy CV. Oral health related quality of life. J Int Soc Prev Community Dent 2013;3(1):1–6.
- **32.** Locker D, Jokovic A, Clarke M. Assessing the responsiveness of measures of oral health-related quality of life. Community Dent Oral Epidemiol 2004;32(1):10–8.

- 33. Copay AG, Subach BR, Glassman SD, Polly Jr DW, Schuler TC. Understanding the minimum clinically important difference: a review of concepts and methods. Spine J 2007;7(5):541–6.
- **34.** Inglehart MR. Oral health and quality of life editors. In: Mostofsky DI, Fortune F, editors. Behavioral dentistry. Hoboken, NJ: Wiley-Blackwell; 2014. p. 11–25.
- Slade GD, Spencer AJ. Development and evaluation of the Oral Health Impact Profile. Community Dent Health 1994;11(1):3– 11.
- **36.** Atchison KA, Dolan TA. Development of the Geriatric Oral Health Assessment Index. J Dent Educ 1990;54(11):680–7.
- Dugas NN, Lawrence HP, Teplitsky P, Friedman S. Quality of life and satisfaction outcomes of endodontic treatment. J Endod 2002;28(12):819–27.
- Slade GD. Derivation and validation of a short-form oral health impact profile. Community Dent Oral Epidemiol 1997;25(4):284–90.
- Montero J, Lorenzo B, Barrios R, Albaladejo A, Miron Canelo JA, Lopez-Valverde A. Patient-centered outcomes of root canal treatment: a cohort follow-up study. J Endod 2015;41(9):1456– 61.
- Liu P, McGrath C, Cheung GS. Improvement in oral healthrelated quality of life after endodontic treatment: a prospective longitudinal study. J Endod 2014;40(6):805–10.
- **41.** Wong MC, Lo EC, McMillan AS. Validation of a Chinese version of the Oral Health Impact Profile (OHIP). Community Dent Oral Epidemiol 2002;30(6):423–30.
- 42. Montero-Martin J, Bravo-Perez M, Albaladejo-Martinez A, Hernandez-Martin LA, Rosel-Gallardo EM. Validation the Oral Health Impact Profile (OHIP-14sp) for adults in Spain. Med Oral Patol Oral Cir Bucal 2009;14(1):E44–50.
- **43.** Szentpétery A, Szabo G, Marada G, Szanto I, John MT. The Hungarian version of the Oral Health Impact Profile. Eur J Oral Sci 2006;114(3):197–203.
- 44. Khoo ST, Ode W, Lopez V, Yu VSH, Lai C, Lui JN. Factors influencing quality of life after surgical and nonsurgical interventions of persistent endodontic disease. J Endod 2020;46 (12):1832–40.
- **45.** McGrath C, Lam O, Lang N. An evidence-based review of patient-reported outcome measures in dental implant research among dentate subjects. J Clin Periodontol 2012;39:193–201.
- 46. Bharathi J, Mittal S, Tewari S, et al. Effect of the piezoelectric device on intraoperative hemorrhage control and quality of life after endodontic microsurgery: a randomized clinical study. J Endod 2021;47(7):1052–60.
- Del Fabbro M, Ceresoli V, Lolato A, Taschieri S. Effect of platelet concentrate on quality of life after periradicular surgery: a randomized clinical study. J Endod 2012;38(6):733–9.
- Del Fabbro M, Taschieri S, Weinstein R. Quality of life after microscopic periradicular surgery using two different incision techniques: a randomized clinical study. Int Endod J 2009;42 (4):360–7.
- 49. Meschi N, Fieuws S, Vanhoenacker A, et al. Root-end surgery with leucocyte- and platelet-rich fibrin and an occlusive membrane: a randomized controlled clinical trial on patients' quality of life. Clin Oral Investig 2018;22(6):2401–11.
- 50. Soto-Peñaloza D, Penarrocha-Diago M, Cervera-Ballester J, Peñarrocha-Diago M, Tarazona-Alvarez B, Peñarrocha-Oltra D. Pain and quality of life after endodontic surgery with or without advanced platelet-rich fibrin membrane application: a randomized clinical trial. Clin Oral Investig 2020;24 (5):1727–38.
- 51. Taschieri S, Corbella S, Tsesis I, Del Fabbro M. Impact of the use of plasma rich in growth factors (PRGF) on the quality of life of patients treated with endodontic surgery when a perforation of sinus membrane occurred. A comparative study. Oral Maxillofac Surg 2014;18(1):43–52.

PROMS IN ENDODONTICS

- 52. Tuk JG, Lindeboom JA, van Wijk AJ. Effect of periapical surgery on oral health-related quality of life in the first postoperative week using the Dutch version of Oral Health Impact Profile-14. Oral Maxillofac Surg 2021;25(4):549–59.
- 53. Tsesis I, Shoshani Y, Givol N, Yahalom R, Fuss Z, Taicher S. Comparison of quality of life after surgical endodontic treatment using two techniques: a prospective study. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2005;99(3):367–71.
- 54. Jordan RA, Markovic L, Holzner AL, Richter B, Gaengler P. Development of a basic root canal treatment (BRT) for primary oral health care–evaluation after one year. Int Dent J 2009;59(3):141–7.
- 55. Yu VS, Messer HH, Yee R, Shen L. Incidence and impact of painful exacerbations in a cohort with post-treatment persistent endodontic lesions. J Endod 2012;38(1):41–6.
- Metin R, Tatli U, Evlice B. Effects of low-level laser therapy on soft and hard tissue healing after endodontic surgery. Lasers Med Sci 2018;33(8):1699–706.
- Wright WG, Jones JA, Spiro 3rd A, Rich SE, Kressin NR. Use of patient self-report oral health outcome measures in assessment of dental treatment outcomes. J Public Health Dent 2009;69(2):95–103.
- 58. Pasqualini D, Corbella S, Alovisi M, et al. Postoperative quality of life following single-visit root canal treatment performed by rotary or reciprocating instrumentation: a randomized clinical trial. Int Endod J 2016;49(11):1030–9.
- 59. Yavari HR, Jafari F, Jamloo H, Hallaj-Nezhadi S, Jafari S. The effect of submucosal injection of corticosteroids on pain perception and quality of life after root canal treatment of teeth with irreversible pulpitis: a randomized clinical trial. J Endod 2019;45(5):477–82.
- 60. Iqbal MS, Rajan S, Iqbal MZ. Determinants of oral healthrelated quality of life among patients on root canal treatment. J Pharm Res Int 2020:76–82.
- Yaylali IE, Teke A, Tunca YM. The effect of foraminal enlargement of necrotic teeth with a continuous rotary system on postoperative pain: a randomized controlled trial. J Endod 2017;43(3):359–63.
- 62. Liu P, McGrath C, Cheung GS. Quality of life and psychological well-being among endodontic patients: a case-control study. Aust Dent J 2012;57(4):493–7.
- 63. Wigsten E, Kvist T, Jonasson P, EndoReCo Davidson T. Comparing quality of life of patients undergoing root canal treatment or tooth extraction. J Endod 2020;46(1):19–28.
- 64. Allen PF, McMillan AS, Walshaw D, Locker D. A comparison of the validity of generic- and disease-specific measures in the assessment of oral health-related quality of life. Community Dent Oral Epidemiol 1999;27(5):344–52.
- **65.** Diniz-de-Figueiredo FE, Lima LF, Oliveira LS, Bernardino IM, Paiva SM, Faria ESAL. The impact of two root canal treatment protocols on the oral health-related quality of life: a randomized controlled pragmatic clinical trial. Int Endod J 2020;53 (53):1327–38.
- 66. Chew T, Brennan D, Rossi-Fedele G. Comparative longitudinal study on the impact root canal treatment and other dental services have on oral health-related quality of life using selfreported health measures (Oral Health Impact Profile-14 and Global Health Measures). J Endod 2019;45(8):985–93 e1.
- 67. He J, White RK, White CA, Schweitzer JL, Woodmansey KF. Clinical and patient-centered outcomes of nonsurgical root canal retreatment in first molars using contemporary techniques. J Endod 2017;43(2):231–7.
- 68. Hamasha AA, Hatiwsh A. Quality of life and satisfaction of patients after nonsurgical primary root canal treatment provided by undergraduate students, graduate students and endodontic specialists. Int Endod J 2013;46(12):1131–9.
- 69. Gatten DL, Riedy CA, Hong SK, Johnson JD, Cohenca N. Quality of life of endodontically treated versus implant treated

patients: a university-based qualitative research study. J Endod 2011;37(7):903–9.

- Michaelson PL, Holland GR. Is pulpitis painful? Int Endod J 2002;35(10):829–32.
- 71. Vena DA, Collie D, Wu H, et al. Prevalence of persistent pain 3 to 5 years post primary root canal therapy and its impact on oral health-related quality of life: PEARL Network findings. J Endod 2014;40(12):1917–21.
- 72. Liu P, McGrath C, Cheung G. What are the key endodontic factors associated with oral health-related quality of life? Int Endod J 2014;47(3):238–45.
- Bartols A, Reutter CA, Robra BP, Walther W. Reciproc vs. hand instrumentation in dental practice: a study in routine care. Peer J 2016;4:e2182.
- 74. Oliveira PS, da Costa KNB, Carvalho CN, Ferreira MC. Impact of root canal preparation performed by ProTaper Next or Reciproc on the quality of life of patients: a randomized clinical trial. Int Endod J 2019;52(2):139–48.
- **75.** Ashari A, Mohamed AM. Relationship of the Dental Aesthetic Index to the oral health-related quality of life. Angle Orthod 2016;86(2):337–42.
- 76. Locker D, Slade G. Association between clinical and subjective indicators of oral health status in an older adult population. Gerodontology 1994;11(2):108–14.
- 77. Guyatt GH, Berman LB, Townsend M, Pugsley SO, Chambers LW. A measure of quality of life for clinical trials in chronic lung disease. Thorax 1987;42(10):773–8.
- Jaeschke R, Singer J, Guyatt GH. Measurement of health status. Ascertaining the minimal clinically important difference. Control Clin Trials 1989;10(4):407–15.
- 79. de Vet HC, Terwee CB, Ostelo RW, Beckerman H, Knol DL, Bouter LM. Minimal changes in health status questionnaires: distinction between minimally detectable change and minimally important change. Health Qual Life Outcomes 2006;4:54.
- **80.** Osoba D, Rodrigues G, Myles J, Zee B, Pater J. Interpreting the significance of changes in health-related quality-of-life scores. J Clin Oncol 1998;16(1):139–44.
- Norman GR, Sloan JA, Wyrwich KW. Interpretation of changes in health-related quality of life: the remarkable universality of half a standard deviation. Med Care 2003;41(5):582–92.
- Engel L, Beaton DE, Touma Z. Minimal clinically important difference: a review of outcome measure score interpretation. Rheum Dis Clin North Am 2018;44(2):177–88.
- Allen PF, O'Sullivan M, Locker D. Determining the minimally important difference for the Oral Health Impact Profile-20. Eur J Oral Sci 2009;117(2):129–34.
- 84. King MT. A point of minimal important difference (MID): a critique of terminology and methods. Expert Rev Pharmacoecon Outcomes Res 2011;11(2):171–84.
- **85.** Mouelhi Y, Jouve E, Castelli C, Gentile S. How is the minimal clinically important difference established in health-related quality of life instruments? Review of anchors and methods. Health Qual Life Outcomes 2020;18(1):136.
- 86. Jayadevappa R, Cook R, Chhatre S. Minimal important difference to infer changes in health-related quality of life-a systematic review. J Clin Epidemiol 2017;89:188–98.
- Revicki D, Hays RD, Cella D, Sloan J. Recommended methods for determining responsiveness and minimally important differences for patient-reported outcomes. J Clin Epidemiol 2008;61(2):102–9.
- Cole JC, Lin P, Rupnow MF. Minimal important differences in the Migraine-Specific Quality of Life Questionnaire (MSQ) version. Cephalalgia 2009;29(11):1180–7.
- 89. Tashjian RZ, Hung M, Keener JD, et al. Determining the minimal clinically important difference for the American Shoulder and Elbow Surgeons score, Simple Shoulder Test, and visual analog scale (VAS) measuring pain after shoulder arthroplasty. J Shoulder Elbow Surg 2017;26(1):144–8.

- **90.** Juniper EF, Guyatt GH, Willan A, Griffith LE. Determining a minimal important change in a disease-specific quality of life questionnaire. J Clin Epidemiol 1994;47(1):81–7.
- **91.** Metz SM, Wyrwich KW, Babu AN, Kroenke K, Tierney WM, Wolinsky FD. A comparison of traditional and Rasch cut points for assessing clinically important change in health-related quality of life among patients with asthma. Qual Life Res 2006;15(10):1639–49.
- **92.** Schunemann HJ, Puhan M, Goldstein R, Jaeschke R, Guyatt GH. Measurement properties and interpretability of the Chronic Respiratory Disease Questionnaire (CRQ). COPD 2005;2(1):81–9.
- Yamazaki M, Inukai M, Baba K, John MT. Japanese version of the Oral Health Impact Profile (OHIP-J). J Oral Rehabil 2007;34 (3):159–68.

- 94. McGrath C, Wong AH, Lo EC, Cheung CS. The sensitivity and responsiveness of an oral health related quality of life measure to tooth whitening. J Dent 2005;33(8):697–702.
- 95. Rener-Sitar K, Petricevic N, Celebic A, Marion L. Psychometric properties of Croatian and Slovenian short form of oral health impact profile questionnaires. Croat Med J 2008;49(4):536–44.
- **96.** Tsakos G, Bernabe E, D'Aiuto F, et al. Assessing the minimally important difference in the oral impact on daily performances index in patients treated for periodontitis. J Clin Periodontol 2010;37(10):903–9.
- **97.** Myint Oo KZ, Fueki K, Yoshida-Kohno E, Hayashi Y, Inamochi Y, Wakabayashi N. Minimal clinically important differences of oral health-related quality of life after removable partial denture treatments. J Dent 2020;92:103246.